


به نام خدا







Classification System





×


STAGES OF ROP












X.

Stage 3



Stage 3 retinopathy of prematurity In stage 3 ROP, new blood vessels and fibrous tissue grow along the ridge and extend into the vitreous.









PLUS

- posterior venous dilation and arteriolar tortuosity of at least 2 quadrants
- Arises gradually or very rapidly.
- Due to AV shunting mainly in ridge tissue
- Severity indicator







PRISTHRESHOLD ROP:

TYPE 1 includes:

- Zone 1- Any ROP with plus ds or stage 3 with or without plus ds
- Zone 2- stage 2 or 3 ROP with plus ds

TYPE 2 includes:

- Zone 1- stage 1 or 2 without plus ds
- Zone 2- stage 3 without plus ds.

Aggressive posterior ROP:

Uncommon,

rapidly progressive,

severe form of ROP,

characterized by its posterior location (usually zone 1) and prominence of plus disease.

Also termed as RUSH disease.

Table 2. Screening Frequency for Patients at Risk for ROP

Timing of Follow-up	Disease
1 week	Zone 1: stage 1 or 2 ROP Zone 2: stage 3 ROP
1-2 weeks	Zone 1: immature Zone 2: stage 2 ROP Zone 1: regressing ROP
2 weeks	Zone 2: stage 1 ROP Zone 2: regressing ROP
2-3 weeks	Zone 2: immature Zone 3: stage 1 or 2 ROP Zone 3: regressing ROP

Screening of at risk babies

Diagnosis

Treatment

- Cryotherapy (mostly outdated)
- · Laser treatment (gold standard)
- Anti-VEGF (adjuvant) before laser and surgery
- Surgery
- Rx of ROP related complications

Post treatment follow up

Rehabilitation



TREATMENT

- RETINAL ABLATION
 - CRYO
 - LASER
- SCLERAL BUCKLING
- VITRECTOMY
 - LENS SPARING
 - With LENSECTOMY






Figure 2 - Schematic drawing showing the moment of application of transscleral diode laser





Thank you for your attention!